Author index to Volume 30, 1993

Key: R = Review
OA = Original article
SR = Short report
CR = Case report
E = Editorial matters
PV = Personal view
L = Letter

Abe F, Miyamoto N, Tahara Y, Takahashi J, Shima K. Serum glycated albumin concentrations during pregnancy, 198 (SR)
Addae S *see* Mohammed S
Adeli K *see* Galvan B
Adzaku F *see* Mohammed S
Aguzzi F, Gasparro C, Bergami MR, Merlini M. High-sensitivity electrophoretic method for the detection of Bence Jones protein and for the study of proteinuria in unconcentrated urines, 287 (OA)
Aiken CGA, Sherwood RA, Lenney W. Role of plasma phosphate measurements in detecting rickets of prematurity and in monitoring treatment, 469 (OA)
Aitken DA *see* Spencer K
Al-Jubouri MA, Young RM. Hyperlipidaemia and diabetes, 331 (L)
Hypertriglyceridaemia: a pointer to diabetes mellitus and alcoholism, 201 (SR)
Al-Saleh IA, Fellows C, Delves T, Taylor A. Identification of sources of lead exposure among children in Arar, Saudi Arabia, 142 (OA)
Albertini A *see* Farina L
Ancarani P *see* Carrea A
Anderson RW *see* Macri JN, and Spencer K
Ang BCN *see* Mazurkiewicz JC
Annobil S *see* Mohammed S
Anthony FW *see* Spencer K
Arendt J *see* English J
Arumanayagam M *see* Chan A
Asquith P *see* Moran A
Athey D, Ball M, McNeil CJ. Avidin-biotin based electrochemical immunoassay for thyrotropin, 570 (OA)

Baadenhuijsen H *see* Jansen RTP
Bailie K *see* Young IS
Baird IM *see* Collinson PO
Ball M *see* Athey D
Ballantyne FC, Gibbons J, O'Reilly DStJ. Urine albumin should replace total protein for the assessment of glomerular proteinuria, 101 (SR)
Barbour H *see* Spencer K
Barbour HM, Logan S. Interaction between bovine control materials and the ferene iron method, 221 (SR)
Barnes DJ *see* Mattock MB
Barnes I *see* Wolff K
Barron JL *see* Madira WM
Beattie GJ, Roulston JE, Eccles DM, Richardson JM, Fisken J, Leonard RCF. Alpha-fucosidase as a marker of genetic deletion in ovarian carcinoma, 207 (OA)
Beattie GJ, Sturgeon C, Fisken J, Roulston JE. The effect of surgery on the serum concentration of CA125 in epithelial ovarian cancer, 402 (SR)
Beetham R, Cattell WR. Proteinuria: pathophysiology, significance and recommendations for measurement in clinical practice, 425 (RA)
Beetham R, Dawnay A, Chany C, Dubrey S, Miles J. A radioimmunoassay for human urinary prealbumin, 377 (OA)
Bell MA *see* Packard CJ
Bergami MR *see* Aguzzi D
Berry E *see* Spencer K
Besser GM *see* Ross GA
Beswick K *see* Hirst AD and Beswick K
Bhatnagar D. The role of clinical biochemistry departments in developing a research infrastructure, 117 (PV)
Bingham SA *see* Mazurkiewicz JC
Blecher M, Sheard AG. Profiting from inventions in academia: American and British perspectives, 1 (PV)
Blumsohn A, Morris B. Is it necessary to alkalinize urine samples for measurement of retinol binding protein? 592 (L)
Bonham JR. The investigation of hypoglycaemia during childhood, 238 (RA)
Bouloux PMG *see* Ross GA
Breimer LH. Would DNA repair assessment benefit patient mangement? 121 (PV)
Broughton PMG *see* Peters M and Broughton PMG
Brown RC *see* Turner G
Bullock DG *see* Jansen RTP
Butler GE *see* Turner G

Canepa A *see* Carrea A
Carrea A, Canepa A, Perfumo F, Ancarani P, Verrina E, Gusmano R. Proteolytic activity and free amino acid concentrations in polymorphonuclear leucocytes, 559 (OA)
Carter GD *see* Collinson PO
Časl M-T, Grubb A. A rapid enzyme-linked immunosorbent assay for serum amyloid A using sequence-specific antibodies, 278 (OA)
Časl M-T, Sabljar-Matovinović M, Kovačević S, Počanić D, Preden-Kereković V, Jagarinec N. Clinical relevance of serum amyloid A protein monitoring in urinary tract infections, 272 (OA)
Cassidy M *see* Packard CJ
Cattell WR *see* Beetham R and Cattell WR
Chan A, Wong F, Arumanayagam M. Serum ultrafiltrable copper, total copper and caeruloplasmin concentrations in gynaecological carcinomas, 545 (OA)
Cheung CK *see* Robertshaw M
Ciechanowicz A, Kawiak J, Miks B, Dutkiewicz T, Dlugosz A, Syczewska-Wawrzynowicz M, Marzecka J. Plasma prekallikrein levels in patients with hepatocellular carcinoma and liver cirrhosis: a pilot study, 445 (OA)
Clark GH, Fraser CG. Biological variation of acute phase proteins, 373 (OA)
Clark PMS *see* Krentz AJ
Cohen MP, Witt J, Wu V-Y. Purified haemoglobin preparations in the evaluation of HbA1c determination by ion exchange chromatography, 265 (OA)

Author index to Volume 30, 1993

Key: R = Review
OA = Original article
SR = Short report
CR = Case report
E = Editorial matters
PV = Personal view
L = Letter

Abe F, Miyamoto N, Tahara Y, Takahashi J, Shima K. Serum glycated albumin concentrations during pregnancy, 198 (SR)
Addae S *see* Mohammed S
Adeli K *see* Galvan B
Adzaku F *see* Mohammed S
Aguzzi F, Gasparro C, Bergami MR, Merlini M. High-sensitivity electrophoretic method for the detection of Bence Jones protein and for the study of proteinuria in unconcentrated urines, 287 (OA)
Aiken CGA, Sherwood RA, Lenney W. Role of plasma phosphate measurements in detecting rickets of prematurity and in monitoring treatment, 469 (OA)
Aitken DA *see* Spencer K
Al-Jubouri MA, Young RM. Hyperlipidaemia and diabetes, 331 (L)
Hypertriglyceridaemia: a pointer to diabetes mellitus and alcoholism, 201 (SR)
Al-Saleh IA, Fellows C, Delves T, Taylor A. Identification of sources of lead exposure among children in Arar, Saudi Arabia, 142 (OA)
Albertini A *see* Farina L
Ancarani P *see* Carrea A
Anderson RW *see* Macri JN, and Spencer K
Ang BCN *see* Mazurkiewicz JC
Annobil S *see* Mohammed S
Anthony FW *see* Spencer K
Arendt J *see* English J
Arumanayagam M *see* Chan A
Asquith P *see* Moran A
Athey D, Ball M, McNeil CJ. Avidin-biotin based electrochemical immunoassay for thyrotropin, 570 (OA)

Baadenhuijsen H *see* Jansen RTP
Bailie K *see* Young IS
Baird IM *see* Collinson PO
Ball M *see* Athey D
Ballantyne FC, Gibbons J, O'Reilly DStJ. Urine albumin should replace total protein for the assessment of glomerular proteinuria, 101 (SR)
Barbour H *see* Spencer K
Barbour HM, Logan S. Interaction between bovine control materials and the ferene iron method, 221 (SR)
Barnes DJ *see* Mattock MB
Barnes I *see* Wolff K
Barron JL *see* Madira WM
Beattie GJ, Roulston JE, Eccles DM, Richardson JM, Fisken J, Leonard RCF. Alpha-fucosidase as a marker of genetic deletion in ovarian carcinoma, 207 (OA)
Beattie GJ, Sturgeon C, Fisken J, Roulston JE. The effect of surgery on the serum concentration of CA125 in epithelial ovarian cancer, 402 (SR)
Beetham R, Cattell WR. Proteinuria: pathophysiology, significance and recommendations for measurement in clinical practice, 425 (RA)
Beetham R, Dawnay A, Chany C, Dubrey S, Miles J. A radioimmunoassay for human urinary prealbumin, 377 (OA)
Bell MA *see* Packard CJ
Bergami MR *see* Aguzzi D
Berry E *see* Spencer K
Besser GM *see* Ross GA
Beswick K *see* Hirst AD and Beswick K
Bhatnagar D. The role of clinical biochemistry departments in developing a research infrastructure, 117 (PV)
Bingham SA *see* Mazurkiewicz JC
Blecher M, Sheard AG. Profiting from inventions in academia: American and British perspectives, 1 (PV)
Blumsohn A, Morris B. Is it necessary to alkalinize urine samples for measurement of retinol binding protein? 592 (L)
Bonham JR. The investigation of hypoglycaemia during childhood, 238 (RA)
Bouloux PMG *see* Ross GA
Breimer LH. Would DNA repair assessment benefit patient mangement? 121 (PV)
Broughton PMG *see* Peters M and Broughton PMG
Brown RC *see* Turner G
Bullock DG *see* Jansen RTP
Butler GE *see* Turner G

Canepa A *see* Carrea A
Carrea A, Canepa A, Perfumo F, Ancarani P, Verrina E, Gusmano R. Proteolytic activity and free amino acid concentrations in polymorphonuclear leucocytes, 559 (OA)
Carter GD *see* Collinson PO
Časl M-T, Grubb A. A rapid enzyme-linked immunosorbent assay for serum amyloid A using sequence-specific antibodies, 278 (OA)
Časl M-T, Sabljar-Matovinović M, Kovačević S, Počanić D, Preden-Kereković V, Jagarinec N. Clinical relevance of serum amyloid A protein monitoring in urinary tract infections, 272 (OA)
Cassidy M *see* Packard CJ
Cattell WR *see* Beetham R and Cattell WR
Chan A, Wong F, Arumanayagam M. Serum ultrafiltrable copper, total copper and caeruloplasmin concentrations in gynaecological carcinomas, 545 (OA)
Cheung CK *see* Robertshaw M
Ciechanowicz A, Kawiak J, Miks B, Dutkiewicz T, Dlugosz A, Syczewska-Wawrzynowicz M, Marzecka J. Plasma prekallikrein levels in patients with hepatocellular carcinoma and liver cirrhosis: a pilot study, 445 (OA)
Clark GH, Fraser CG. Biological variation of acute phase proteins, 373 (OA)
Clark PMS *see* Krentz AJ
Cohen MP, Witt J, Wu V-Y. Purified haemoglobin preparations in the evaluation of HbA1c determination by ion exchange chromatography, 265 (OA)

Colbert DL, Turner GJ, Gooch JC. Reducing the cost of phenytoin assays with in-house TDx™ reagents, 555 (OA)
Cole M. Measurement of dopamine, HVA and HMMA in untimed urine samples, 499 (L)
Collinson PO, Moseley D, Stubbs PJ, Carter GD. Troponin T for the differential diagnosis of ischaemic myocardial damage, 11 (OA)
Collinson PO, Ramhamadamy EM, Stubbs PJ, Rosalki SB, Garrat HM, Mosely D, Evans DH, Fink RS, Baird IM, Greenwood TW. Rapid enzyme diagnosis of patients with acute chest pain reduces patient stay in the coronary care unit, 17 (OA)
Collinson PO *see also* Johnston JD
Cook EJ *see* Macri JN
Coombes EJ *see* Spencer K
Cooreman W *see* Uyttenbroeck W
Cowdrey GN, Tasker PJ, Gould BJ, Rice-Oxley M, Firth GB. Isoelectric focusing in an immobilized pH gradient for the detection of intrathecal IgG in cerebrospinal fluid: sensitivity and specificity for the diagnosis of multiple sclerosis, 463 (OA)
Cox L *see* Krentz AJ
Creagh FN *see* Turner G
Crook M. Hyponatraemia and hypouricaemia, 217 (SR)
Crook M *see also* Haq M
Crossley JA *see* Spencer K
Csathy L *see* Price RG

Dagen MM *see* Packard CJ
Davie SJ, Whiting KL, Gould BJ. Biological variation in glycated proteins, 260 (OA)
Davies GG *see* Morris RHK and Davies GG
Davies PTG *see* Hughes PJ
Dawnay A *see* Beetham R
De Leenheer A *see* Jansen RTP
de Verdier C-H *see* Jansen RTP
Dean JJ *see* Killeen AA
Dear PRF *see* Henderson MJ and Dear PRF
Delves T *see* Al-Saleh IA
Demacker PNM *see* Janssens PMW
Dick JPR *see* Hughes PJ
Dlugosz A *see* Ciechanowicz A
Dominiczak MH *see* Kilpatrick ES
Donders SHJ, Lustermans FAT, van Wersch JWJ. The effect of microalbuminuria on glycaemic control, serum lipids and haemostasis parameters in non-insulin-dependent diabetes mellitus, 439 (OA)
Donovan CM *see* Wright DE
Doran J *see* Spencer K
Dornan TL *see* Taylor AJ and Dornan TL
Dubrey S *see* Beetham R
Duffy F *see* Knopf H-P
Duggan PF *see* O'Leary N
Dumont G *see* Jansen RTP
Dutkiewicz T *see* Ciechanowicz A

Eastwood MA *see* Fowlie S
Eaton RH *see* Packard CJ
Eccles DM *see* Beattie GJ
Edwards GA *see* Graham PE
Ellis AR. Antenatal screening for Down's syndrome—can we do better? 421 (PV), 590 (Addendum)
English J, Middleton BA, Arendt J, Wirz-Justice A. Rapid direct measurement of melatonin in saliva using an iodinated tracer and solid phase second antibody, 415 (SR)
Ernerudh J *see* Öhman S
Evans DH *see* Collinson PO

Fairly I *see* Robertshaw M
Farina L, Iacobello C, Orlandini S, Ius A, Albertini A. The effect of the concentration ratio of avidin and biotin on a single step sandwich enzyme immunoassay, 87 (OA)
Fell GS *see* Martin BJ
Fellows C *see* Al-Saleh IA
Fielding AM. Diagnosis of familial benign hypercalcaemia, 111 (L)
Fink RS *see* Collinson PO
Firth GB *see* Cowdrey GN
Fisher RM, Potts JL, Frayn KN. Immunoturbidimetric measurement of low concentrations of apolipoprotein B in lipaemic samples, 584 (SR)
Fisken J *see* Beattie GJ
Foo AY *see* Thum CN
Forsberg P *see* Öhman S
Fowlie S, Eastwood MA, Wenham PR. Salivary immunoglobulin A in irritable bowel syndrome: no correlation with anxiety, 324 (SR)
Fraser CG *see* Clark GH and Fraser CG
Fraser WD *see* Manning EMC and Fraser WD
Frayn KN *see* Fisher RM
Freaney R *see* McKenna MJ and Freaney R
Fry IDR *see* Starkey BJ and Fry IDR

Galvan B, Macri J, Adeli K. An amplified enzyme-linked immunosorbent assay for rapid and sensitive determination of apolipoprotein AI in plasma and tissue culture media, 404 (SR)
Garrat HM *see* Collinson PO
Gasparro C *see* Aguzzi D
Ghany C *see* Beetham R
Gibbons J *see* Ballantyne FC
Gillett MG *see* Gunneberg A
Glen ACA *see* Kilpatrick WS
Goldie DJ *see* Gunneberg A
Gooch JC *see* Colbert DL
Gould BJ *see* Cowdrey GN, Davie SJ, and Kirkham PA
Graham PE, Smythe GA, Edwards GA, Lazarus L. Laboratory diagnosis of phaeochromocytoma: which analytes should we measure? 129 (OA); 506 (L)
Greenwood TW *see* Collinson PO
Grossman A *see* Ross GA
Grubb A *see* Casl M-T and Grubb A
Gunn IR, Wallace JR. Diagnosis of familial benign hypercalcaemia, 111 (L)
Gunneberg A, Gillett MG, Goldie DJ. A high serum concentration of azlocillin affects routine biochemical analyses, 329 (CR)
Gusmano R *see* Carrea A
Guy JM, McMurray JR. Urinary retinol binding protein: stability and pre-analytical handling of specimens for its measurement, 77 (OA); 592 (L)

Hackshaw A *see* Wald N
Haddow JE *see* Wald N

Handley G, Johnson PG, Peel D. Raised serum fructosamine concentration caused by IgA paraproteinaemia, 106 (CR)
Haq M, Haq S, Tutt P, Crook M. Serum total sialic acid and lipid-associated sialic acid in normal individuals and patients with myocardial infarction, and their relationship to acute phase proteins, 383 (OA)
Haq S *see* Haq M
Harrison SP. Delay in separation of blood samples, 108 (L)
Hay AWM *see* Wolff K
Hayashi Y *see* Matsumoto S
Henderson AR. Assessing test accuracy and its clinical consequences: a primer for receiver operating characteristic curve analysis, 521 (RA)
Henderson MJ. Measurement of dopamine, HVA and HMMA in untimed urine samples, 499 (L)
Henderson MJ, Dear PRF. Role of the clinical biochemistry laboratory in the management of very low birth-weight infants, 341 (RA); erratum, 462
Heys AD *see* Trewick AL
Hillis A *see* Marsden JT
Hirst AD, Beswick K. A blood spot assay for apo A1 and B lipoproteins and the apo B/A1 ratio, 476 (OA)
Hughes PJ, Kiely PDW, Davies PTG, Dick JPR, Turner AM, Lane RJM. Hyperammonaemia—an important cause of encephalopathy: three case reports, 225 (CR)
Hutchinson E *see* Nisbet JA

Iacobello C *see* Farina L
Ijlst L, Wanders RJA. A simple spectrophotometric assay for long-chain acyl-CoA dehydrogenase activity measurements in human skin fibroblasts, 293 (OA)
Ikata T *see* Matsumoto S
Ius A *see* Farina L
Izatt D. Glycated haemoglobin, 590 (L)

Jagarinec N *see* Časl M-T
James TJ *see* Nurse DR and James TJ
Jansen RTP, Bullock DG, Vassault A, Baadenhuijsen H, De Leenheer A, Dumont G, de Verdier C-H, Zender R. Between-country comparability of clinical chemistry results: an international quality assessment survey of 17 analytes in six European countries through existing national schemes, 304 (OA)
Janssens PMW, Vonk J, Demacker PNM. Hypertriglyceridaemia in a case of subcutaneous fat necrosis in a newborn, 482 (SR)
Jennings A *see* Satur CMR
Johnson PG *see* Handley G
Johnston J, McLelland A, O'Reilly DStJ. The relationship between serum cholesterol and serum thyroid hormones in male patients with suspected hypothyroidism, 256 (OA)
Johnston JD, Collinson PO, Rosalki SB. MIDAS: Myocardial infarct diagnosis by assessment of slope, 407 (SR)
Johnston JD, Koneru S, Kuwana T, Rosalki SB. Bone-derived serum enzymes and bone density in perimenopausal Caucasian women, 191 (OA)
Jones AF *see* Moran A
Jones GRD. HemoCue blood glucose analyser and haematocrit, 510 (L)
Jung K, Priem F. What are the criteria to introduce new methods for determination of urinary N-acetyl-β-D-glucosaminidase activity? 501 (L)

Kaddoumi O *see* Mohammed S
Kawiak J *see* Ciechanowicz A
Keen H *see* Mattock MB
Kemp GJ. Hypophosphataemia and leukaemia, 590 (L)
Kemp PM, McGrann RG, Sampson WFD. Mononuclear blood cell magnesium, 507 (L)
Kenny D. Method comparison, 501 (L)
Kiely PDW *see* Hughes PJ
Killeen AA, Ramey ML, Dean JJ. High-dose hook effect in an immunoluminometric thyrotropin assay: the open-faced sandwich artefact, 413 (SR)
Kilpatrick ES, Rumley AG, Myint H, Dominiczak MH, Small M. The effect of variations in haematocrit, mean cell volume and red blood cell count on reagent strip tests for glucose, 485 (SR)
Kilpatrick WS, Wosornu D, McGuiness JB, Glen ACA. Early diagnosis of acute myocardial infarction: CK-MB and myoglobin compared, 435 (OA)
Kimura S *see* Yamada T
Kinoshita I *see* Matsumoto S
Kirkham PA, Petry CJ, Morris BA, Gould BJ. Glycated 6-aminohexanoic acid - an improved calibrator for the serum fructosamine assay, 410 (SR)
Knight GJ *see* Wald N
Knopf H-P, Duffy F , Papoian R. A time-resolved fluoroimmunoassay for recombinant human interleukin-3, 69 (OA)
Komori T *see* Yoshimura M
Koneru S *see* Johnston JD
Kovacevic S *see* Casl M-T
Krentz AJ, Clark PMS, Cox L, Williams AC, Nattrass M. Insulin and proinsulin-like molecules in motor neurone disease, 195 (SR)
Kuwana T *see* Johnston JD
Kyne F *see* McGing PG and Kyne F
Maguire S and Kyne F

Labib MH *see* Palfrey SM and Labib MH
Lane RJM *see* Hughes PJ
Lazarus L *see* Graham PE
Lee LT, Maldonado BR, Roheim PS, Thompson JJ. Analysis of apolipoprotein size distribution by electroimmunoblotting from non-denaturing composite gels, 157 (OA)
Lenney W *see* Aiken CGA
Leonard RCF *see* Beattie GJ
Leslie H *see* Stevenson HP
Lester E *see* West P
Logan S *see* Barbour HM and Logan S
Lustermans FAT *see* Donders SHJ
Luxton F *see* Trewick AL
Lyon TDB *see* Martin BJ

McGing PG, Kyne F. Cutbacks in cardiac enzyme testing, 418 (L)
McGrann RG *see* Kemp PM
McGuiness JB *see* Kilpatrick WS
McKenna MJ, Freaney R. Secondary hyperparathyroidism, 504 (L)
McLelland A *see* Johnston J
McMurray JR *see* Guy JM and McMurray JR
McNeil CJ *see* Athey D

Macri J *see* Galvan B
Macri JN, Spencer K, Anderson RW, Cook EJ. Free beta-chorionic gonadotropin: a cross-reactivity study of two immunometric assays used in prenatal maternal serum screening for Down's syndrome, 94 (OA)
see also Spencer K
Madira WM, Wilcox AH, Barron JL. Storage of plasma in primary plasma separator tubes, 213 (SR)
Maguire S, Kyne F. A rapid selective high-performance liquid chromatography assay for cyclosporine, 488 (SR)
Maldonado BR *see* Lee LT
Male IA *see* Salzmann MB and Male IA
Mallard AS *see* Spencer K
Mallios RR. Predicting the probability of helper T cell immunodominant sites through discriminant analysis, 152 (OA)
Manning EMC, Fraser WD. A survey of diagnoses in patients with a low intact parathyroid hormone concentration, 252 (OA)
Marsden JT, Sherwood RA, Hillis A, Peters TJ. Monitoring erythropoietin therapy for anaemia of chronic renal failure by serum erythropoietin assays, 205 (SR)
Martin BJ, Lyon TDB, Walker W, Fell GS. Mononuclear blood cell magnesium, 507 (L)
Mononuclear blood cell magnesium in older subjects: evaluation of its use in clinical practice, 23 (OA)
Marzecka J *see* Ciechanowicz A
Massey J *see* Page K and Massey J
Masters PW. Audit of automated requesting of serum magnesium measurement in patients with hypocalcaemia, 248 (OA)
Matsumoto S, Hayashi Y, Kinoshita I, Ikata T, Yamamoto S. Immunoaffinity purification of prostaglandin E2 and leukotriene C4 prior to radioimmunoassay: application to human synovial fluid, 60 (OA)
Matsuto T *see* Yamada T
Mattock MB, Barnes DJ, Keen H. Screening for microalbuminuria in a mixed ethnic diabetic clinic, 509 (L)
Mazurkiewicz J *see* Nisbet JA
Mazurkiewicz JC, Bingham SA, Runswick S, Ang BCN. Inhibition of the urease reaction by boric acid, 215 (SR)
Merlini M *see* Aguzzi D
Middleton BA *see* English J
Miks B *see* Ciechanowicz A
Miles J *see* Beetham R
Milford Ward A *see* Spencer K
Miller AL. Plasma bicarbonate assays - time for a new look? 233 (PV)
Miyamoto N *see* Abe F
Mohammed S, Addae S, Suleiman S, Adzaku F, Annobil S, Kaddoumi O, Richards J. Serum calcium, parathyroid hormone, and vitamin D status in children and young adults with sickle cell disease, 45 (OA)
Moran A, Radley S, Neoptolemos J, Jones AF, Asquith P. Detection of colorectal cancer by faecal alpha-1-antitrypsin, 28 (OA)
Morris B *see* Blumsohn A and Morris B
Morris BA *see* Kirkham PA
Morris RHK, Davies GG. Thin layer chromatographic identification of thromboxane B2 in serum, and platelet rich plasma stimulated by arachidonic acid and collagen, 203 (SR)
Moseley D *see* Collinson PO
Mosely D *see* Collinson PO
Muskiet FAJ *see* Weykamp CW
Myint H *see* Kilpatrick ES

Nakagawa T *see* Ueda K
Nakajima H *see* Ueda K
Nakanishi T, Shimizu A. Identification of 3-hydroxyisovalerylcarnitine in the urine of a patient with multiple carboxylase deficiency, 318 (SR)
see also Yoshimura M
Nattrass M *see* Krentz AJ
Neoptolemos J *see* Moran A
Newbould EC *see* Ross GA
Nisbet JA, Hutchinson E, Mazurkiewicz J. Europium labelled time-resolved fluoroimmunoassay for urine albumin, 223 (SR)
Nomata Y *see* Yamada T
Nurse DR, James TJ. An adaptable computer interface for radioimmunoassay, 298 (OA)

O'Brien PA *see* Tormey WP and O'Brien PA
O'Donnell J *see* Smellie WSA
Oelbaum RS *see* Thum CN
Ogunleye IO, Olusi SO. Purification and catalytic properties of liver alcohol dehydrogenase isoenzymes in patients with hepatoma in Nigeria, 146 (OA)
Öhman S. Intrathecally produced IgG, 417 (L), 590 (Errata)
Öhman S, Ernerudh J, Forsberg P, von Schenck H, Vrethem M. Improved formulae for the judgement of intrathecally produced IgA and IgM in the presence of blood CSF barrier damage, 454 (OA)
Okada M *see* Yamada T
Olah VA *see* Price RG
O'Leary N, Pembroke A, Duggan PF. A robust procedure for the automated measurement of total serum bilirubin using potassium ferricyanide, 175 (OA)
Olusi SO *see* Ogunleye IO and Olusi SO
O'Reilly DStJ *see* Ballantyne FC, and Johnston J
Orlandini S *see* Farina L

Packard CJ, Bell MA, Eaton RH, Dagen MM, Cassidy M, Shepherd J. A pilot scheme for improving the accuracy of serum cholesterol measurement in Scotland and Northern Ireland, 387 (OA)
Page K, Massey J. Urine catecholamine excretion in long-term oral theophylline use, 492 (SR)
Palfrey SM, Labib MH. A simple method for measuring neopterin in serum using HPLC, 209 (SR)
Palomaki GE *see* Wald N
Papoian R *see* Knopf H-P
Patel S *see* Trewick AL
Peel D *see* Handley G
Pembroke A *see* O'Leary N
Penders TJ *see* Weykamp CW
Perfumo F *see* Carrea A
Perrett D *see* Ross GA
Peters M, Broughton PMG. The role of expert systems in improving the test requesting patterns of clinicians, 52 (OA)

Peters TJ *see* Marsden JT
Petry CJ *see* Kirkham PA
Piscator, 115, 232, 336, 420, 520, 602
Počanić D *see* Casl M-T
Pocsi I *see* Price RG
Potts JL *see* Fisher RM
Preden-Kereković V *see* Časl M-T
Price CP. Multiple forms of human serum alkaline phosphatase: detection and quantitation, 355 (RA)
Price RG, Richardson AC, Pocsi I, Csathy L, Olah VA. What are the criteria to introduce new methods for determination of urinary N-acetyl-β-D-glucosaminidase activity? 501 (L)
Priem F *see* Jung K and Priem F

Radley S *see* Moran A
Ramey ML *see* Killeen AA
Ramhamadamy EM *see* Collinson PO
Reynolds TM *see* Wright DE
Rice-Oxley M *see* Cowdrey GN
Richards J *see* Mohammed S
Richardson AC *see* Price RG
Richardson JM *see* Beattie GJ
Riches P, Rippin J, Walker S. Seminal fluid contamination of urine, 228 (CR)
Rippin J *see* Riches P
Roberts NB *see* Tewari A
Robertshaw M, Cheung CK, Fairly I, Swaminathan R. Protein excretion after prolonged exercise, 34 (OA)
Robertson WS. Laboratory diagnosis of phaeochromocytoma, 506 (L)
Roheim PS *see* Lee LT
Rosalki SB *see* Collinson PO, Johnston JD, and Thum CN
Ross GA, Newbould EC, Thomas J, Bouloux PMG, Besser GM, Perrett D, Grossman A. Plasma and 24 h-urinary catecholamine concentrations in normal and patient populations, 38 (OA)
Roulston JE *see* Beattie GJ
Rumley AG *see* Kilpatrick ES
Runswick S *see* Mazurkiewicz JC

Sabljar-Matovinovic M *see* Casl M-T
Salzmann MB, Male IA. Keep the lid on it: artefactual hypernatraemia in samples from paediatric patients, 211 (SR)
Sampson WFD *see* Kemp PM
Sanderson MJ *see* Wolff K
Sandler D *see* Smellie WSA
Satur CMR, Jennings A, Walker DR. Hypomagnesaemia and fits complicating paediatric cardiac surgery, 315 (SR)
Savory DJ, Whitlock T. Serum creatinine: effect of delay in separation, 508 (L)
Scharpe S *see* Uyttenbroeck W
Schuller E. Intrathecally produced IgG, 417 (L)
Sells RA *see* Tewari A
Sheard AG *see* Biecher M and Sheard AG
Shepherd J *see* Packard CJ
Sheridan B *see* Stevenson HP
Sherwood RA *see* Aiken CGA, and Marsden JT
Shima K *see* Abe F
Shimizu A *see* Nakanishi T and Shimizu A, and Ueda K
Simpson WG. Delay in separation of blood samples, 108 (L)
Skinner A *see* Turner G
Small M *see* Kilpatrick ES
Smellie WSA, Sandler D, O'Donnell J. Hyperlipidaemia and diabetes, 331 (L)
Smythe GA *see* Graham PE
Spencer K, Coombes EJ, Mallard AS, Milford Ward A. Use of free beta-hCG in Down's syndrome screening, 512 (L)
Spencer K, Macri JN, Anderson RW, Aitken DA, Berry E, Crossley JA, Wood PJ, Coombes EJ, Stroud M, Worthington DJ, Doran J, Barbour H, Wilmot R. Dual analyte immunoassay in neural tube defect and Down's syndrome screening: results of a multicentre clinical trial, 394 (OA)
Spencer K, Wood PJ, Anthony FW. Elevated levels of maternal serum inhibin immunoreactivity in second trimester pregnancies affected by Down's syndrome, 219 (SR)
Spencer K *see also* Macri JN
Starkey BJ, Fry IDR. Comparison of a commercial enzymic kit for urinary oxalate analysis with a high performance liquid chromatographic method, 186 (OA)
Stevenson HP, Leslie H, Sheridan B. Serum free beta-human chorionic gonadotrophin concentrations increase in unseparated blood specimens, 99 (SR)
Stroud M *see* Spencer K
Stubbs PJ *see* Collinson PO
Sturgeon C *see* Beattie GJ
Sugita O *see* Yamada T
Suleiman S *see* Mohammed S
Swaminathan R. Parathyroid hormone measurement, 109 (L)
see also Robertshaw M
Syczewska-Wawrzynowicz M *see* Ciechanowicz A

Tahara Y *see* Abe F
Takahashi H *see* Yoshimura M
Takahashi J *see* Abe F
Tasker PJ *see* Cowdrey GN
Taylor A *see* Al-Saleh IA
Taylor AJ, Dornan TL. Successful treatment of short stature and delayed puberty in congenital magnesium-losing kidney, 494 (CR)
Tewari A, Ward RG, Sells RA, Roberts NB. Reduced bioavailability of cyclosporin A capsules in a renal transplant patient with partial gastrectomy and ileal resection, 587 (CR)
Thomas J *see* Ross GA
Thompson JJ *see* Lee LT
Thum CN, Oelbaum RS, Foo AY, Rosalki SB. Renal isoamylase clearance as a measure of altered renal charge selectivity in patients with diabetes mellitus, 449 (OA)
Tindall H *see* West P
Tormey WP, O'Brien PA. Clinical associations of an increased transthyretin band in routine serum and urine protein electrophoresis, 550 (OA)
Trewick AL, Luxton F, Patel S, Heys AD. Urine protein electrophoresis without sample concentration: an evaluation of two methods, 321 (SR)
Trimble ER *see* Young IS
Turner AM *see* Hughes PJ

Turner G, Brown RC, Weeks I, Butler GE, Creagh FN, Woodhead JS. Urinary growth hormone excretion as measured by a sensitive immunochemiluminometric assay, 180 (OA)
Turner G, Skinner A, Woodhead JS. Urinary growth hormone measurements in children with renal insufficiency, 540 (OA)
Turner GJ *see* Colbert DL
Tutt P *see* Haq M

Ueda K, Nakajima H, Nakagawa T, Shimizu A. The association between hepatitis C virus infection and *in vitro* activation of the complement system, 565 (OA)
Uyttenbroeck W, Cooreman W, Scharpe S. The presence of antibodies against extractable nuclear antigens in serum: a comparison of immunoblotting versus radial immunodiffusion, 83 (OA)

Vadgama P. Biosensors in clinical biochemistry, 337 (PV)
van der Slik W *see* Weykamp CW
van Wersch JWJ *see* Donders SHJ
Vasikaran SD. Index of individuality and utility of population based reference range, 500 (L), 590 (Errata), 594 (L)
Vassault A *see* Jansen RTP
Verrina E *see* Carrea A
von Schenck H *see* Öhman S
Vonk J *see* Janssens PMW
Vrethem M *see* Öhman S

Wald N, Hackshaw A, Haddow JE, Palomaki GE, Knight GJ. Use of free beta-hCG in Down's syndrome screening, 512 (L)
Walker DR *see* Satur CMR
Walker S *see* Riches P
Walker W *see* Martin BJ
Wallace JR *see* Gunn IR and Wallace JR
Wanders RJA *see* Ijlst L and Wanders RJA
Ward RG *see* Tewari A
Weeks I *see* Turner G
Wenham PR. Molecular biology techniques, 332 (L)
Wenham PR *see also* Fowlie S
West P, Tindall H, Lester E. Screening for microalbuminuria in a mixed ethnic diabetic clinic, 104 (SR)
Weykamp CW, Penders TJ, Muskiet FAJ, van der Slik W. Glycohaemoglobin: comparison of 12 analytical methods, applied to lyophilized haemolysates by 101 laboratories in an external quality assurance programme, 169 (OA)
Whiting KL *see* Davie SJ
Whitlock DJ *see* Savory DJ and Whitlock T
Wiener K. An assessment of the effect of haematocrit on the HemoCue blood glucose analyser, 90 (OA); 510 (L)
Wierzbicki AS. Molecular biology techniques, 332 (L)
Wilcox AH *see* Madira WM
Williams AC *see* Krentz AJ
Wilmot R *see* Spencer K
Wirz-Justice A *see* English J
Witt J *see* Cohen MP
Wolff K, Sanderson MJ, Hay AWM, Barnes I. An evaluation of the measurement of drugs of abuse by commercial and in-house horizontal thin layer chromatography, 163 (OA)
Wong F *see* Chan A
Wood PJ *see* Spencer K
Woodhead JS. Parathyroid hormone measurement, 109 (L)
Woodhead JS *see also* Turner G
Worthington DJ *see* Spencer K
Wosornu D *see* Kilpatrick WS
Wright DE, Reynolds TM, Donovan CM. Assessment of atypicality: an adjunct to prenatal screening for Down's syndrome that facilitates detection of other abnormalities, 578 (OA)
Wu V-Y *see* Cohen MP

Yamada T, Nomata Y, Sugita O, Okada M. A rapid method for measuring serum amyloid A protein by latex agglutination nephelometric immunoassay, 72 (OA)
Yamada T, Sugita O, Okada M, Matsuto T, Kimura S. Use of a chemiluminescent acridinium ester label for cellular uptake assay of low density lipoprotein, 490 (SR)
Yamamoto S *see* Matsumoto S
Yoshimura M, Komori T, Nakanishi T, Takahashi H. Estimation of sulphoconjugated catecholamine concentrations in plasma by high-performance liquid chromatography, 135 (OA)
Young IS, Bailie K, Trimble ER. Severe hypophosphataemia in a patient with acute leukaemia, 326 (CR); 590 (L)
Young RM *see* Al-Jubouri MA and Young RM

Zender R *see* Jansen RTP

Subject index to Volume 30, 1993

Key: R = Review; OA = Original article; SR = Short report; CR = Case report; E = Editorial matters; PV = Personal view; L = Letter

Acute phase proteins
biological variation, 373 (OA)
and serum total and lipid-associated sialic acid, in normal patients and myocardial infarction, 383 (OA)
Albumin
glycated, serum concentrations during pregnancy, 198 (SR)
urine, europium labelled time-resolved fluoroimmunoassay, 223 (SR)
urine, should replace total protein for assessment of glomerular proteinuria, 101 (SR)
Albuminuria after prolonged exercise, 34 (OA)
Alcoholism, hypertriglyceridaemia as pointer, 201 (SR)
Alkaline phosphatase, serum, multiple forms, detection and quantitation, 355 (RA)
Alpha-1-antitrypsin, faecal, in detection of colorectal cancer, 28 (OA)
Alpha-fucosidase as marker of genetic deletion in ovarian carcinoma, 207 (SR)
6-Aminohexanoic acid, glycated, improved calibrator for serum fructosamine assay, 410 (SR)
Amniocentesis, adjunct to Down's syndrome screening that facilitates detecion of other abnormalities, 578 (OA)
Amyloid A protein, serum
ELISA using sequence-specific antibodies, 278 (OA)
measurement by latex agglutination nephelometric immunoassay, rapid method, 72 (OA)
monitoring, clinical relevance in urinary tract infections, 272 (OA)
Anaemia of chronic renal failure, erythropoietin therapy, monitoring by serum erythropoietin assays, 205 (SR)
Antenatal screening, for Down's screening, can we do better? 421 (PV), 590 (Addendum)
Antibodies, against extractable nuclear antigens, immunoblotting and radial immunodiffusion compared, 83 (OA)
Apolipoprotein
A1, amplified ELISA for rapid and sensitive determination, 404 (SR)
A1 and B lipoproteins, blood spot assay, and apo B/A1 ratio, 476 (OA)
B, immunoturbidimetric measurement of low concentrations in lipaemic samples, 584 (SR)
size distribution, analysis by electroimmunoblotting from non-denaturing composite gels, 157 (OA)
Arachidonic acid, -stimulated platelet rich plasma, thin layer chromatographic identification of thromboxane B2, 203 (SR)
Assessing test accuracy and its clinical consequences, primer for ROC curve analysis, 521 (RA)
Audit for magnesium measurement for hypocalcaemia, 248 (OA)
Avidin:biotin concentration, effect on single step sandwich enzyme immunoassay, 87 (OA)
Avidin-biotin based electrochemical immunoassay for thyrotropin, 570 (OA)
Azlocillin, high serum concentrations affect routine biochemical analyses, 329 (CR)

Bartter's syndrome, successful treatment for short stature and delayed puberty, 494 (CR)
Bayesian discriminant analysis to predict probability of helper T cell immunodominant sites, 152 (OA)
Bence Jones protein
electrophoresis using unconcentrated urine, 287 (OA)
electrophoresis without sample concentration, 321 (SR)
Between-country comparability of clinical chemistry results, international quality assessment survey of 17 analytes in six European countries through existing national schemes, 304 (OA)
Bicarbonate, plasma assays, time for a new look? 233 (PV)
Bilirubin, total serum, robust procedure for automated measurement using potassium ferricyanide, 175 (OA)
Biosensors in clinical biochemistry, 337 (PV)
Biotin:avidin concentration, effect on single step sandwich enzyme immunoassay, 87 (OA)
Biotin-avidin based electrochemical immunoassay for thyrotropin, 570 (OA)
Blood samples, delay in separation, 108 (L); 508 (L)
Blood spot assay for A1 and B lipoproteins, and apo B/A1 ratio, 476 (OA)
Bloom's syndrome, DNA repair assessment, 121 (PV)
Bone-derived serum enzymes and bone density, in perimenopausal Caucasian women, 191 (OA)
Book reviews
Allen HJ, Kisailus EC, *editors*. Glycoconjugates—composition, structure, and function, 1992, 597
Bierich JR, *editor*. Clinical endocrinology and metabolism: international practice and research. Vol 6, No. 3, Growth disorders, 1992, 598
Fernandes J, Saudubray JM, Tada K, *editors*. Inborn metabolic diseases. Diagnosis and treatment, 1990, 231
Flatt PR, *editor*. Nutrient regulation of insulin secretion, 1992, 597
Kricka LJ. Nonisotopic DNA probe techniques, 1992, 598
Krstulovic AM, *editor*. Chiral separations by HPLC—applications to pharmaceutical compounds, 335
Nakamura RM, *editor*. Immunochemical assays and biosensor technology of the 1990s, 1992, 114
Plebani M, Di Mario F. Advances in gastroenterology. Gastric secretion: biochemical and clinical features, 1992, 599
Rifai N, Warwick GR, *editors*. Methods for clinical chemistry measurement of lipid and lipoprotein risk factors, 1991, 596
Soldin SJ, Rifai N, Hicks JMB, *editors*. Biochemical basis of pediatric disease, 1992, 519
Young DS. Effects of drugs on clinic chemistry laboratory tests—1991 supplement to third edition, 1991, 596

Boric acid, in inhibition of urease reaction, 215 (SR)
Bovine control materials and ferene iron method, interaction between, 221 (SR)

CA125, serum concentration in epithelial ovarian cancer, effect of surgery, 402 (SR)
Calcium, serum, in sickle cell disease in children and young adults, 45 (OA)
Carcinomas, gynaecological, serum ultrafiltrable copper, total copper and caeruloplasmin concentrations, 545 (OA)
Cardiac enzyme testing, cutbacks, 418 (L)
Cardiac surgery, paediatric, hypomagnesaemia and fits complicating, 315 (SR)
Catecholamine concentrations
plasma and 24h-urinary, normal and patient populations, 38 (OA)
sulphoconjugated, estimation in plasma by HPLC, 135 (OA)
urine excretion in long-term oral theophylline use, 492 (SR)
Centrifugation and artefactual hypernatraemia in samples from paediatric patients, 211 (SR)
Cerebrospinal fluid
blood barrier damage, improved formulae for judgement of intrathecally produced IgA and IgM, 454 (OA)
intrathecal IgG, in multiple sclerosis, isoelectric focusing, 463 (OA)
Cervix, carcinoma, serum ultrafiltrable copper, total copper and caeruloplasmin concentrations, 545 (OA)
Chemiluminescent acridinium ester label, use for cellular uptake assay of low density lipoprotein, 490 (SR)
Chest pain, acute, rapid enzyme diagnosis reduces patient stay in coronary care unit, 17 (OA)
Cholesterol
serum, measurement, pilot scheme for improving accuracy, Scotland and Northern Ireland, 387 (OA)
and thyroid hormones, serum, relationship in suspected hypothyroidism, males, 256 (OA)
Clinical biochemistry departments, role in developing a research infrastructure, 117 (PV)
Clinical biochemistry laboratory, role in management of very low birthweight infants, 341 (RA); erratum, 462
Collagen-stimulated platelet rich plasma, thin layer chromatographic identification of thromboxane B2, 203 (SR)
Colorectal cancer, detection by faecal alpha-1-antitrypsin, 28 (OA)
Complement system activation, *in vitro*, and HCV infection, association between, 565 (OA)
Copper, serum, ultrafiltrable and total, and caeruloplasmin concentrations, in gynaecological carcinomas, 545 (OA)
Creatine kinase increase rate in acute chest pain, and stay in coronary care unit, 17 (OA)
Creatine kinase MB and myoglobin compared, for early diagnosis of myocardial infarction, 435 (OA)
Creatinine, serum, effect of delay in separation, 508 (L)
Cyclosporin A capsules, reduced bioavailability in renal transplant patient with partial gastrectomy and ileal resection, 587 (CR)
Cyclosporine, rapid selective HPLC assay, 488 (SR)

Diabetes mellitus
and hyperlipidaemia, 331 (L)
hypertriglyceridaemia as pointer, 201 (SR)
renal isoamylase clearance as measure of altered renal charge selectivity, 449 (OA)
and serum glycated albumin concentrations during pregnancy, 198 (SR)
Diabetic clinic, mixed ethnic, microalbuminuria screening, 104 (SR); 509 (L)
DNA-based techniques, diagnostic application, 332 (L)
DNA repair assessment, benefit in patient management, 121 (PV)
Dopamine, measurement in untimed urine samples, 499 (L)
Down's syndrome
adjunct to prenatal screening facilitating detection of other abnormalities, 578 (OA)
affecting second trimester pregnancies, and elevated levels of maternal serum inhibin immunoreactivity, 219 (SR)
antenatal screening, can we do better? 421 (PV), 590 (Addendum)
and neural tube defect screening, dual analyte immunoassay, 394 (OA)
prenatal screening, free beta-hCG, cross-reactivity of two assays, 94 (OA)
use of free beta-hCG in screening, 512 (L)
Drugs of abuse, evaluation of measurement by commercial and in-house horizontal thin layer chromatography, 163 (OA)
Dual analyte immunoassay in neural tube defect and Down's syndrome screening, 394 (OA)

Electrochemical immunoassay, avidin-biotin based, for thyrotropin, 570 (OA)
Electroimmunoblotting from non-denaturing composite gels, in analysis of apolipoprotein size distribution, 157 (OA)
Electrolytes, plasma bicarbonate assays, time for a new look? 233 (PV)
Electrophoresis
of Bence Jones protein, in unconcentrated urine, 287 (OA)
increased transthyretin band, clinical associations, 550 (OA)
urine protein, without sample concentration, 321 (SR)
Encephalopathy, hyperammonaemia an important cause, 225 (CR)
Enzyme-linked immunosorbent assay
amplified, for rapid and sensitive determination of apoAI, 404 (SR)
for serum amyloid A, using sequence-specific antibodies, 278 (OA)
Epithelial ovarian cancer, effect of surgery on serum concentration of CA125, 402 (SR)
Erratum, 462
Erythropoietin assays for monitoring erythropoietin therapy for anaemia of chronic renal failure, 205 (SR)
Erythropoietin therapy, for anaemia of chronic renal failure, monitoring by serum erythropoietin assays, 205 (SR)
Ethnic minority groups, screening for microalbuminuria in diabetic clinic, 104 (SR); 509 (L)
Europium labelled time-resolved fluoroimmunoassay, for urine albumin, 223 (SR)

Exercise, prolonged, protein excretion after, 34 (OA)
Expert systems, role in improving test requesting of clinicians, 52 (OA)
External quality assessment, between-country comparability, 304 (OA)
External quality assurance programme, comparison of 12 analytical methods, 169 (OA)

Familial benign hypercalcaemia, diagnosis, 111 (L)
Ferene iron method and bovine control materials, interaction between, 221 (SR)
Fibroblasts, skin, long-chain acyl-CoA dehydrogenase activity measurements, spectrophotometry, 293 (OA)
Fits, and hypomagnesaemia, complicating paediatric cardiac surgery, 315 (SR)
Fluoroimmunoassay
europium labelled time-resolved, for urine albumin, 223 (SR)
time-resolved, for recombinant human interleukin-3, 69 (OA)
Free amino acid concentrations and proteolytic activity, in polymorphonuclear leucocytes, 559 (OA)
Fructosamine
raised serum concentration caused by IgA paraproteinaemia, 106 (CR)
serum, assay, glycated 6-aminohexanoic acid an improved calibrator, 410 (SR)

Glomerular proteinuria, urine albumin should replace total protein for assessment, 101 (SR)
Glucose, reagent strip tests, effect of variations in haematocrit, mean cell volume and red blood cell count, 485 (SR)
Glycated 6-aminohexanoic acid, improved calibrator for serum fructosamine assay, 410 (SR)
Glycated albumin, serum concentrations during pregnancy, 198 (SR)
Glycated haemoglobin, 590 (L)
in determination of HbA1c by ion exchange chromatography, 265 (OA)
Glycated proteins, biological variation, 260 (OA)
Glycohaemoglobin, external quality assurance programme, comparison of 12 analytical methods, 169 (OA)
Growth hormone
urinary excretion as measured by sensitive immunochemiluminometric assay, 180 (OA)
urinary measurements in children with renal insufficiency, 540 (OA)

Haematocrit
effect on HemoCue blood glucose analyser, 90 (OA); 510 (L)
mean cell volume and red blood cell count, variations, effect on reagent strip tests for glucose, 485 (SR)
Haemoglobin, purified preparations, in HbA1c determination by ion exchange chromatography, 265 (OA)
Helper T cell immunodominant sites, predicting probability through discriminant analysis, 152 (OA)
HemoCue blood glucose analyser, effect of haematocrit, 90 (OA); 510 (L)
Hepatitis C virus infection, and *in vitro* activation of complement system, association between, 565 (OA)
Hepatocellular carcinoma, and liver cirrhosis, plasma prekallikrein levels, pilot study, 445 (OA)
Hepatoma, purification and catalytic properties of liver alcohol dehydrogenase isoenzymes, Nigeria, 146 (OA)
High-performance liquid chromatography
and commercial enzymic kit compared, for urinary oxalate analysis, 186 (OA)
in estimation of sulphoconjugated catecholamine concentrations in plasma, 135 (OA)
for measuring neopterin in serum, 209 (SR)
rapid selective, for cyclosporine, 488 (SR)
Homovanillic acid, measurement in untimed urine samples, 499 (L)
Horizontal thin layer chromatography, commercial and in-house, in evaluation of measurement of drugs of abuse, 163 (OA)
Human chorionic gonadotrophin, free beta
concentrations increase in unseparated blood specimens, 99 (SR)
cross-reactivity study of two assays, prenatal Down's syndrome screening, 94 (OA)
in Down's syndrome screening, 512 (L)
3-Hydroxyisovalerylcarnitine in urine, multiple carboxylase deficiency, 318 (SR)
Hydroxymethylmandelic acid, measurement in untimed urine samples, 499 (L)
Hyperammonaemia, important cause of encephalopathy, 225 (CR)
Hypercalcaemia, familial benign, diagnosis, 111 (L)
Hyperlipidaemia, and diabetes, 331 (L)
Hypernatraemia, artefactual, in samples from paediatric patients, 211 (SR)
Hyperparathyroidism, secondary, 504 (L)
Hypertriglyceridaemia
pointer to diabetes mellitus and alcoholism, 201 (SR)
in subcutaneous fat necrosis in a newborn, 482 (SR)
Hypocalcaemia, serum magnesium measurement, audit, 248 (OA)
Hypoglycaemia, during childhood, investigation, 238 (RA)
Hypomagnesaemia and fits, complicating paediatric cardiac surgery, 315 (SR)
Hyponatraemia, and hypouricaemia, study of plasma urate concentrations, 217 (SR)
Hypophosphataemia, severe, in acute leukaemia, 326 (CR), 590 (L)
Hypothyroidism, relationship between serum cholesterol and thyroid hormones, males, 256 (OA)
Hypouricaemia, and hyponatraemia, study of plasma urate concentrations, 217 (SR)

Immunoblotting and radial immunodiffusion compared, in presence of antibodies against extractable nuclear antigens, 83 (OA)
Immunochemiluminometric assay, sensitive, measuring urinary growth hormone excretion, 180 (OA)
Immunoglobulin A
paraproteinaemia, causing raised serum fructosamine concentration, 106 (CR)
salivary, in irritable bowel syndrome, no correlation with anxiety, 324 (SR)
Immunoglobulin A and M, intrathecally produced, and blood CSF barrier damage, improved formulae of judgement, 454 (OA)
Immunoglobulin G
intrathecal, in CSF, isoelectric focusing, and diagnosis of MS, 463 (OA)
intrathecally produced, 417 (L), 590 (Errata)

Immunoturbidimetric measurement of low concentrations of apo B in lipaemic samples, 584 (SR)
Inhibin immunoreactivity, maternal serum, elevated levels in second trimester pregnancies affected by Down's syndrome, 219 (SR)
Insulin and proinsulin-like molecules, in motor neurone disease, 195 (SR)
Interleukin-3, recombinant human, time-resolved fluoroimmunoassay, 69 (OA)
Inventions, in academia, profiting from, American and British perspectives, 1 (PV)
Ion exchange chromatography, purified haemoglobin preparations in evaluation of HbA1c determination, 265 (OA)
Irritable bowel syndrome, salivary IgA, no correlation with anxiety, 324 (SR)
Ischaemic myocardial damage, troponin T for differential diagnosis, 11 (OA)
Isoamylase, renal clearance as measure of altered renal charge selectivity in diabetes mellitus, 449 (OA)

Laboratory investigations, patterns of clinicians' request, role of expert systems, 52 (OA)
Laboratory results, between-country comparability, 304 (OA)
Latex agglutination nephelometric immunoassay, rapid method for measuring serum amyloid A protein, 72 (OA)
Lead exposure, identification of sources among children in Arar, Saudi Arabia, 142 (OA)
Leukaemia, acute, and severe hypophosphataemia, 326 (CR), 590 (L)
Leukotriene C4, and prostaglandin E2, immunoaffinity purification prior to radioimmunoassay, synovial fluid, 60 (OA)
Lipaemic samples, immunoturbidimetric measurement of low concentrations of apo B, 584 (SR)
Liver alcohol dehydrogenase isoenzymes, purification and catalytic properties in hepatoma, Nigeria, 146 (OA)
Liver cirrhosis and hepatocellular carcinoma, plasma prekallikrein levels, pilot study, 445 (OA)
Long-chain acyl-CoA dehydrogenase activity measurements in skin fibroblasts, spectrophotometry, 293 (OA)
Low density lipoprotein, use of chemiluminescent acridinium ester label for cellular uptake assay, 490 (SR)

Magnesium
-losing kidney, congenital, successful treatment of short stature and delayed puberty, 494 (CR)
measurement, serum, for hypocalcaemia, audit, 248 (OA)
mononuclear blood cell, in older subjects, evaluation of its use in clinical practice, 23 (OA); 507 (L)
Melatonin, rapid direct measurement in saliva using iodinated tracer and solid phase second antibody, 415 (SR)
Method comparison studies, 501 (L)
Microalbuminuria
effect on glycaemic control, serum lipids and haemostasis parameters in NIDDM, 439 (OA)
screening in mixed ethnic diabetic clinic, 104 (SR); 509 (L)
MIDAS, myocardial infarct diagnosis by assessment of slope, 407 (SR)
Molecular biology techniques, 332 (L)
Mononuclear blood cell magnesium in older subjects, evaluation of its use in clinical practice, 23 (OA); 507 (L)
Motor neurone disease, insulin and proinsulin-like molecules, 195 (SR)
Multiple carboxylase deficiency, 3-hydroxyisovalerylcarnitine in urine, 318 (SR)
Multiple sclerosis
intrathecal IgA and IgM in blood CSF barrier damage, 454 (OA)
intrathecal IgG in CSF, isoelectric focusing, 463 (OA)
Myocardial infarction
diagnosis by assessment of slope (MIDAS), 407 (SR)
early diagnosis, CK-MB and myoglobin compared, 435 (OA)
early enzyme exclusion reduces patient stay in coronary care unit, 17 (OA)
serum total and lipid-associated sialic acid, and acute phase proteins, 383 (OA)
troponin T for differential diagnosis, 11 (OA)
Myoglobin and CK-MB compared, for early diagnosis of myocardial infarction, 435 (OA)

N-acetyl-β-D-glucosaminidase, what are criteria to introduce new methods for determination of urinary activity, 501 (L)
Neopterin, measurement in serum using HPLC, 209 (SR)
Neural tube defect and Down's syndrome screening, dual analyte immunoassay, 394 (OA)
Non-insulin-dependent diabetes mellitus, effect of microalbuminuria on glycaemic control, serum lipids and haemostasis parameters, 439 (OA)

Osteoporosis, bone-derived serum enzymes and bone density in perimenopausal Caucasian women, 191 (OA)
Ovarian carcinoma
alpha-fucosidase as marker of genetic deletion, 207 (SR)
serum ultrafiltrable copper, total copper and caeruloplasmin concentrations, 545 (OA)
Oxalate, urinary analysis, HPLC and commercial enzymic kit compared, 186 (OA)

Paediatric cardiac surgery, hypomagnesaemia and fits complicating, 315 (SR)
Parathyroid hormone
concentration, low intact, diagnoses, 252 (OA)
measurement, 109 (L), 504 (L)
in sickle cell disease in children and young adults, 45 (OA)
Phaeochromocytoma, laboratory diagnosis, which analytes should we measure? 129 (OA); 506 (L)
Phenytoin assays, reducing cost with in-house TDx™ reagents, 550 (OA)
Phosphate, plasma measurements in detecting rickets of prematurity and in monitoring treatment, 469 (OA)
Plasma, storage in primary plasma separator tubes, 213 (SR)
Population based reference range, index of individuality and utility, 500 (L), 590 (Errata), 594 (L)
Polymerase chain reaction technique, diagnostic application, 332 (L)
Polymorphonuclear leucocytes, proteolytic activity and free amino acid concentrations, 559 (OA)
Potassium ferricyanide, robust procedure for automated measurement of total serum bilirubin, 175 (OA)

Prealbumin, human urinary, radioimmunoassay, 377 (OA)
Pregnancy, serum glycated albumin concentrations during, 198 (SR)
Prekallikrein plasma levels in hepatocellular carcinoma and liver cirrhosis, pilot study, 445 (OA)
Profiting from inventions in academia, American and British perspectives, 1 (PV)
Prostaglandin E2, and leukotriene C4, immunoaffinity purification prior to radioimmunoassay, synovial fluid, 60 (OA)
Protein excretion after prolonged exercise, 34 (OA)
Proteinuria
glomerular, urine albumin should replace total protein for assessment, 101 (SR)
pathophysiology, significance and recommendations for measurement in clinical practice, 425 (RA)
Proteolytic activity and free amino acid concentrations in polymorphonuclear leucocytes, 559 (OA)

Radial immunodiffusion and immunoblotting compared, in presence of antibodies against extractable nuclear antigens, 83 (OA)
Radioimmunoassay
adaptable computer interface, 298 (OA)
for human urinary prealbumin, 377 (OA)
Reagent strip tests for glucose, effect of variations in haematocrit, mean cell volume and red blood cell count, 485 (SR)
Receiver operating characteristic curve analysis, assessing test accuracy and clinical consequences, 521 (RA)
Renal charge selectivity, altered, in diabetes mellitus, renal isoamylase clearance as measure, 449 (OA)
Renal failure, chronic, anaemia, erythropoietin therapy, monitoring by serum erythropoietin assays, 205 (SR)
Renal insufficiency, urinary growth hormone measurements, children, 540 (OA)
Renal transplant patient with partial gastrectomy and ileal resection, and reduced bioavailability of CyA, 587 (CR)
Research infrastructure, role of clinical biochemistry departments, 117 (PV)
Research profits, American and British perspectives, 1 (PV)
Retinol binding protein, urinary, stability and preanalytical handling of specimens for its measurement, 77 (OA), 592 (L)
Rickets of prematurity, plasma phosphate measurements in detecting and in monitoring treatment, 469 (OA)

Salivary immunoglobulin A in irritable bowel syndrome, no correlation with anxiety, 324 (SR)
Screening, for microalbuminuria, in mixed ethnic diabetic clinic, 104 (SR); 509 (L)
Seminal fluid contamination of urine, 228 (CR)
Separation of blood samples, delay, 108 (L); 508 (L)
Sequence-specific antibodies, in ELISA for serum amyloid A, 278 (OA)
Sialic acid, total serum, and lipid-associated, normal and in myocardial infarction, and acute phase proteins, 383 (OA)
Sickle cell disease, serum calcium, parathyroid hormone, and vitamin D status, children and young adults, 45 (OA)
Single step sandwich enzyme immunoassay, effect of avidin:biotin concentration, 87 (OA)
Spectrophotometry, for long-chain acyl-CoA dehydrogenase activity measurements in skin fibroblasts, 293 (OA)
Statistical techniques, method comparison studies, 501 (L)
Subcutaneous fat necrosis in a newborn, and hypertriglyceridaemia, 482 (SR)
Synovial fluid, immunoaffinity purification of PGE2 and leukotriene C4 prior to radioimmunoassay, 60 (OA)

TDx™ reagents, in-house, in reducing cost of phenytoin assays, 550 (OA)
Test accuracy, assessing, and its clinical consequences, primer for ROC curve analysis, 521 (RA)
Test requesting patterns of clinicians, expert systems, role, 52 (OA)
Theophylline, oral, long-term, urine catecholamine excretion, 492 (SR)
Therapeutic drug monitoring, reducing costs with in-house TDx™ reagents, 550 (OA)
Thin layer chromatographic identification of thromboxane B2 in serum, and platelet rich plasma stimulated by arachidonic acid and collagen, 203 (SR)
Thromboxane B2, in serum, and platelet rich plasma stimulated by arachidonic acid and collagen, thin layer chromatographic identification, 203 (SR)
Thyroid and cholesterol hormones, serum, relationship in suspected hypothyroidism, males, 256 (OA)
Thyrotropin
avidin-biotin based electrochemical immunoassay, 570 (OA)
immunoluminometric assay, high-dose hook effect, open-faced sandwich artefact, 413 (SR)
Transthyretin, increased band in routine electrophoresis, clinical associations, 550 (OA)
Troponin T, for differential diagnosis of ischaemic myocardial damage, 11 (OA)

Urease reaction, inhibition by boric acid, 215 (SR)
Urinary tract infections, serum amyloid A protein monitoring, clinical relevance, 272 (OA)
Urine
protein electrophoresis, without sample concentration, 321 (SR)
seminal fluid contamination, 226 (CR)

Very low birthweight infants, role of clinical biochemistry laboratory in management, 341 (RA); erratum, 462
Vitamin D status, in sickle cell disease in children and young adults, 45 (OA)

Xeroderma pigmentosum, DNA repair assessment, 121 (PV)